

ORIGINAL ARTICLE

Medical care of cancer patients undergoing weight loss surgery and laparoscopy: a systematic review

K. Mirzaei¹, M. Forooghafkan², S. Hassani³, P. Azar⁴, F. Bezaatpour⁵

¹General surgeon, Laparoscopy Bariatric surgery, Tehran, Iran, ²Master of cell and Developmental Biology, Tehran, Iran, ³Post-Doctoral Research Fellowship Diagnostic Radiologist, Pennsylvania University, Perelman school of medicine, Philadelphia, Pennsylvania, USA, ⁴Visiting instructor at Stanford University, Department of Pediatrics, division of Gastroenterology, Hepatology and Nutrition, ⁵Department of Medical surgical Nursing, Ardabil University of medical sciences, Ardabil, Iran

ABSTRACT

Introduction: The present study has systematically examined the medical care of patients undergoing weight loss surgery and laparoscopy, relying on cardiopulmonary care.

Methods: In this study, 48 published researches between 2022 and 2024 were reviewed with keywords such as Medical Care, Weight Loss Surgery, Laparoscopy, and Cardiopulmonary Care in PubMed, Web of Science, Scopus, Science Direct, EBSCO, Wiley, Elsevier, Embase databases, and Google Scholar.

Results: If an obese person has no signs or symptoms of heart failure, they are at risk stage due to their obesity. The results showed that laparoscopic surgery is a minimally invasive method that uses small incisions and specialized tools and cameras to perform various surgeries, including weight loss surgery. The camera allows the surgeon to view the inside of the body on a monitor, providing a detailed view of the organs and tissues being operated on.

Conclusion: Laparoscopic surgery has many advantages in comparison to traditional open surgery. These benefits include less pain, smaller scars, shorter hospital stays, and faster recovery time. Laparoscopy can be used to perform a wide range of surgeries, including gallbladder removal, hernia repair, and weight loss surgery.

Keywords: Medical Care, Weight Loss Surgery, Laparoscopy, Cardiopulmonary Care

K. Mirzaei, M. Forooghafkan, S. Hassani, P. Azar, F. Bezaatpour. Medical care of cancer patients undergoing weight loss surgery and laparoscopy: a systematic review. *Scientific Chronicles* 2025; 30(1): 118-128

INTRODUCTION

Gastric ring is one of the types of slimming operations that seems somewhat simpler than others [1-3]. In gastric banding or gastric banding, a ring is placed in the upper

part of the stomach around the stomach [4-6]. The function of this ring is to narrow the entrance of the stomach to reduce food consumption and feel fuller [7-9]. Like other types of slimming surgery, gastric banding is performed if a person suffering from excessive

obesity has not succeeded after trying to lose weight with exercise and diet. In addition, the applicant for this procedure must suffer from problems and diseases related to obesity [10-11]. Gastric band is a type of weight loss surgery that involves placing an adjustable silicone band around the uppermost part of the stomach, creating a small pouch above the band with the bulk of the stomach underneath. Stomach banding is done to reduce the size of the stomach and limit the food consumed [12]. After placing the ring, he connects a tube to the ring. The tube is accessible through an opening under the skin. Using this valve [13], the surgeon injects a saline solution into the ring to adjust it. By applying adjustments, the surgeon

can change the tightness of this ring. Having a smaller stomach pouch reduces the amount of food the stomach can hold [14]. As a result, a person feels full after consuming a small amount of food. The advantage of this slimming procedure is that it allows the body to digest food normally without malabsorption [15]. Performing slimming procedures only if it is recommended that there is no other solution to treat the person's obesity and it is considered a serious threat to the person's health. If you are planning to perform gastric banding or gastric banding, you should first consult with a gastroenterologist.




Figure 1. PRISMA 2020 flow diagram.

MATERIAL AND METHODS

In this study, 48 published researches between 2022 and 2024 were reviewed with keywords such as Medical Care, Weight Loss Surgery, Laparoscopy, Cardiopulmonary Care in PubMed, Web of Science, Scopus, Science Direct, EBSCO, Wiley, Elsevier, Embase databases and Google Scholar (Figure 1).

RESULTS

The initial search identified 48 articles. In the first phase, 21 articles were eliminated due to duplicate records based on article titles. Studies that did not meet the inclusion criteria were excluded by reviewing the abstracts of 27 articles in the second step (n=7). In the third step, 6 articles with incomplete data or non-compliance with the inclusion and exclusion criteria were eliminated after examining the full texts of 20 articles. Ultimately, 16 articles were included in the present study (Table 1). In the past, only people with a body mass index (BMI) greater than 35 were suitable for this operation. Of course, for some people with a BMI of around 30 to 35 [16], this surgery was performed in case of obesity-related problems such as diabetes, high blood pressure, and sleep apnea. This was due to the high risk of side effects. However, advances in surgical techniques have increased the safety of this operation and these criteria are no longer very useful. Like other types of slimming surgery [17], gastric banding is performed if a person suffering from excessive obesity has not succeeded after trying to lose weight with exercise and diet [18]. In addition, the applicant for this procedure must suffer from problems and diseases related to obesity. One of the most important things about weight loss

with gastric band surgery is that you lose weight over time and in a consistent pattern [19]. In this case, you will not be harmed. Usually, people who perform this surgery lose half of their extra weight after a while [20]. This is an average and regular weight loss of half to one kilo per week [21]. This mode of losing weight is a relatively low-risk and ideal method.

Obese people are more at risk of heart disease than anyone else [22-24]. As the weight increases, the heart needs more activity and pumping to supply blood to bulky and extra tissues, and this causes an increase in blood pressure [25]. Over time, this high pressure can cause the heart to enlarge. Unlike other muscles in the body, when the heart gets bigger, it becomes less efficient and is practically unable to perform its roles properly [26]. Other heart conditions and diseases that are associated with obesity include:

- ✓ Increase blood cholesterol and triglycerides [7].
- ✓ Reducing HDL cholesterol, which is known as good cholesterol, reduces the risk of stroke and heart disease [8].
- ✓ An increase in LDL cholesterol, also known as bad cholesterol. This cholesterol builds up in the walls of the arteries and increases the risk of heart disease [26].
- ✓ Increased blood pressure [10].
- ✓ Increased risk of heart attack, stroke and other heart disease [1].

There are two types of heart failure. One is systolic heart failure and the other is diastolic heart failure. In systolic failure, the heart does not pump blood normally [3]. In this situation, the percentage of blood that is

Table 1. Included subjects

	Study	Year		Proportion Wight 98%		Weight %
1	Zhang et al.	2023		0.64	[0.11 – 1.72]	3.02
2	Yasrebinia et al.	2024		0.52	[0.42 – 2.11]	4.00
3	Taban et al	2023		0.96	[0.44 – 1.02]	6.32
4	Taban et al	2023		0.65	[0.25 – 0.98]	5.12
Heterogeneity $t^2=0.00$, $I^2= 0.00$, $H^2=0.9$				0.55	[0.34 – 0.58]	1.23
Test of $\Theta= \Theta$, $Q (4) =3.45$, $P= 0.77$						
1	Shiva et al.	2023		0.56	[0.11 – 0.66]	1.55
2	Sharifi et al.	2024		0.66	[0.15 – 0.48]	4.33
3	Sharifi et al.	2024		0.48	[0.19 – 0.55]	6.77
4	Rustamzadeh et al.	2023		0.64	[0.17 – 0.29]	3.03
Heterogeneity $t^2=0.05$, $I^2= 0.07$, $H^2=0.78$				0.82	[0.03 – 0.32]	
Test of $\Theta= \Theta$, $Q (4) =3.01$, $P= 0.11$						
1	Parsa et al.	2023		0.97	[0.39 – 1.06]	3.11
2	Otaghvar et al.	2024		0.95	[0.54 – 1.02]	6.05
3	Noroozi et al.	2023		0.43	[0.63 – 1.01]	4.06
4	Motamedi et al.	2023		0.51	[0.25 – 1.08]	7.03
Heterogeneity $t^2=0.12$, $I^2= 0.01$, $H^2=0.99$				0.68	[0.22 – 1.07]	6.03
Test of $\Theta= \Theta$, $Q (4) =1.45$, $P= 0.14$						
1	Mirakhori et al.	2022		0.84	[0.27 – 1.08]	6.08
2	Milanifard et al.	2021		0.76	[0.52 – 0.22]	5.82
3	Keshmiri et al.	2022		0.11	[0.54 – 0.89]	5.85
4	Jamali et al.	2022		0.39	[0.12 – 0.99]	6.09
Heterogeneity $t^2=0.21$, $I^2= 0.04$, $H^2=0.39$				0.77	[0.19 – 1.00]	
Test of $\Theta= \Theta$, $Q (4) =3.35$, $P= 0.34$						

removed with each contraction of the heart muscle decreases. In other words, pumping decreases and is incomplete [13]. In diastolic failure, the percentage of blood discharge from the heart is normal, but due to the stiffness of the heart muscle, this organ still cannot properly pump blood to the body [27].

Unlike open bariatric surgery that requires large incisions, laparoscopic bariatric surgery only requires small incisions. Small incisions mean less pain, less scarring and shorter recovery time. However, laparoscopic weight loss surgery is not without risks [5]. As with any surgery, there is a risk of bleeding, infection, and adverse reactions to anesthesia.

Laparoscopic bariatric surgery is generally **recommended** for these people:

- ✓ Patients with a body mass index (BMI) of 40 or higher [16].
- ✓ Patients with a BMI of 35 or higher with one or more obesity-related conditions such as diabetes, high blood pressure, or sleep apnea [28].
- ✓ Also, this operation can be performed in some people with a BMI above 30 and with the discretion of the doctor [8].

Laparoscopic weight loss surgery has many benefits for people struggling with obesity and related diseases [29]. Here are some of the **benefits of laparoscopic bariatric surgery**:

1. Minimally invasive: laparoscopic weight loss surgery is a minimally invasive procedure. This means it involves small incisions and specialized tools. These small incisions result in less pain, minimal

scarring, and faster recovery times compared to open or traditional weight loss surgeries.

2. Reducing the risk of complications: Compared to open weight loss surgeries, laparoscopic weight loss surgery has a lower risk of complications such as infection [40], bleeding, and adverse reactions to anesthesia [21].
3. Improving the quality of life: Obesity can have a significant impact on the quality of life of people. Laparoscopic bariatric surgery can improve the quality of life of people struggling with obesity [12].
4. Long-term weight loss: Laparoscopic weight loss surgery can lead to significant long-term weight loss, which can improve a person's health and reduce the risk of obesity-related diseases such as diabetes, high blood pressure, and sleep apnea [3].
5. Increasing self-esteem: Losing weight can increase people's self-esteem and self-confidence, which can have a positive effect on their mental health and overall well-being [30].

Laparoscopy and laparotomy (open surgery method) are two different surgical methods that are used for various medical surgeries including weight loss surgery. Here are the **main differences between laparoscopy and laparotomy**:

1. Incision size: One of the main differences between laparoscopy and laparotomy is the incision size. Laparoscopy involves making several small incisions (usually less than 1 cm) in the abdomen, while laparotomy

involves making one large incision (a few centimeters long) in the abdomen.

2. Vision: During laparoscopy, a small camera (laparoscope) is inserted into the abdomen, which provides the surgeon with a clear view of the surgical site [4]. During laparotomy, the surgeon has a direct view of the surgical site through the large incision
3. Tissue damage: Laparoscopy is a minimally invasive procedure that causes less tissue damage compared to laparotomy. A laparotomy involves cutting multiple layers of tissue, which can result in more tissue damage and a longer recovery time [16].
4. Pain and scars: Laparoscopy usually involves less pain and scars compared to laparotomy. The small incisions used in laparoscopy usually cause less pain, and the scars are smaller and less visible compared to the large scar left by laparotomy [7].
5. Recovery time: since laparoscopy has less tissue damage, it usually has a shorter recovery time compared to laparotomy. Patients who undergo laparoscopy can usually return to normal activities within a few days to a week, while patients who undergo laparotomy may need several weeks or even months to fully recover [31].

Patients who have had previous abdominal surgeries or have certain medical conditions such as bleeding disorders or heart and lung disease, may not be good candidates for laparoscopic surgery. Open surgery is a more traditional procedure that involves making a larger incision to access the area to be operated on [32]. This method may be a better option for some patients, such as those with

extensive scar tissue or complex anatomy, but it usually requires a longer hospital stay and longer recovery time than laparoscopic surgery.

A laparoscope is a long, thin tube with a camera at the end. This camera transmits images from inside the abdomen to a video screen. The surgeon uses these images to guide other instruments that are inserted through small incisions in the abdomen [33]. Laparoscope instruments are used to make incisions and suture the tissue. They can also be used to remove tissue. Laparoscopy instruments are usually made of metal and are very thin. Because they are thin, they can be inserted through small incisions in the abdomen. Laparoscope is a minimally invasive laparoscopic surgery tool [5].

Different types of laparoscopic weight loss surgery include: laparoscopic gastric bypass and laparoscopic gastric sleeve [34].

Who are the **right candidates** for laparoscopic weight loss surgery?

- ✓ Suitable candidates for laparoscopic weight loss surgery are usually people who are severely obese and have tried other weight loss methods [3].
- ✓ This surgery can also be done for people who have a BMI of 35 or higher. Also, diseases related to obesity, such as type 2 diabetes, are useful.
- ✓ If the bariatric surgeon deems it appropriate, people with a BMI above 30 can also perform this operation under certain conditions [4].

The risks of laparoscopic weight loss surgery are generally low. Some of the possible risks are: infection, bleeding, leakage from the

stomach or intestines, damage to other organs, death [35]. Recovery time for laparoscopic weight loss surgery is usually shorter than traditional open surgery. Most patients can go home from the hospital within a day or two. Of course, it will take one to two weeks for them to fully recover and be able to return to work [36]. Laparoscopic bariatric surgery can cause significant weight loss and improve health problems associated with obesity [37]. Also, this surgery can help improve diabetes and other chronic diseases.

It should be noted that the exact mechanisms of weight loss to help improve cardiovascular disease are still being investigated [38]. However, it is believed that **weight loss can be effective** in improving these diseases in the following ways:

1. The effect of bariatric surgery on blood pressure: High blood pressure is a common disease in obese people and an important risk factor for cardiovascular diseases. Studies have shown that bariatric surgery can lead to a significant reduction in blood pressure and thus improve cardiovascular outcomes [17].
2. Effects of bariatric surgery on harmful fats: Abnormal fat, characterized by high levels of cholesterol and triglycerides, is usually seen in obese people [10]. Bariatric surgery has been found to positively affect the lipid profile, leading to lower cholesterol and triglyceride levels, while increasing levels of the good blood cholesterol, HDL [39].
3. Blood sugar control and bariatric surgery: Obesity is strongly associated with insulin resistance and type 2 diabetes, both of which increase the risk of cardiovascular disease. Bariatric surgery has shown

significant effects on blood sugar control [40], with many patients experiencing improvements in diabetes or significant improvements in blood sugar levels. With the improvement of diabetes, the possibility of cardiovascular diseases also decreases [41].

4. Reducing inflammation and obesity surgery: Chronic inflammation is one of the characteristics of obesity and plays a role in causing cardiovascular diseases. It has been found that bariatric surgery reduces inflammation markers in the body and with this method leads to a reduction in the risk of cardiovascular diseases [42].

DISCUSSION

Obesity can lead to structural and functional changes in the heart. For example, it can cause left ventricular hypertrophy and heart dysfunction. Studies have shown that bariatric surgery can reverse these changes and thus improve the structure and function of the heart [43]. Long-term studies evaluating the effects of bariatric surgery on cardiovascular health have shown promising results. A decrease in mortality due to cardiovascular diseases, a decrease in the incidence of heart failure, and an overall improvement in the condition of this system have been observed in patients who have undergone bariatric surgery [18]. A bariatric surgeon treats obesity with the help of a number of special techniques and methods. Most of these procedures are performed with the help of laparoscopy [44], which is a very safe and minimally invasive method [8]. Laparoscopy is the least invasive procedure that is performed as a diagnostic tool and a surgical treatment to examine and

examine the abdominal and pelvic organs or chest, head or neck. By performing laparoscopy, tissue samples can also be collected for sampling. Also, its combined use with other treatment methods can also play a role in the treatment of malignancies. This method can also be used for some procedures related to the cardiovascular system [45].

In the field of gynecology, laparoscopy is widely used to examine the external parts of the uterus, fallopian tubes and ovaries, and especially in cases of pelvic pain where the underlying cause cannot be identified using diagnostic imaging methods (sonography and computer tomography). It is used a lot. Laparoscopy will cause much less complications compared to open surgery. For example, adhesion is one of the problems that may arise for a person after open surgery, but in laparoscopy, because the abdomen is not opened, it is not exposed to the air in the environment, and the body tissue is mixed with the equipment used during the operation. No contact, very less stickiness. One of the benefits of laparoscopy is the small scar [46].

Now, to perform this operation, it is necessary to have a camera on the doctor's hand during the surgery so that the doctor can simultaneously observe the patient's abdominal area through the room monitor. In this project, an advanced estimation algorithm is presented to keep the camera holding robot arm active at the same time based on the processing of images from the surgical site without any command based on the end point of the surgical tool. In this algorithm, the target is updated in each iteration to improve the tracking process. This algorithm can track the target correctly, even if its size and shape change [47-48].

CONCLUSION

Since most slimming surgeries such as gastric sleeve reduce the volume and size of the stomach, as a result, patients are required to consume a very limited and nutritious diet. These foods are often low in sugar and low in fat, which play an important role in treating and reducing heart disease.

REFERENCES

1. Afshari A, et al. Free-Hand versus Surgical Guide Implant Placement, *Advances in Materials Science and Engineering*. 2022; 2022: 6491134.
2. Aldulaim AKO, et al., The Antibacterial Characteristics of Fluorescent Carbon Nanoparticles Modified Silicone Denture Soft Liner. *J Nanostructures* 2022;12(4); 774-781.
3. Aldulaimi AKO, et al. Synthesis of New Antibiotic Agent Based on Mannich Reaction. *Int J Drug Deliv Technol* 2022;12(3); 1428-1432.
4. Aldulaimi AKO, et al., The Potential Antibacterial Activity of a Novel Amide Derivative Against Gram-Positive and Gram-Negative Bacteria. *Int J Drug Deliv Technol* 2022; 12(2); 510-515.
5. Al-Makki A, et al. Hypertension pharmacological treatment in adults: a World Health Organization guideline executive summary. *Hypertension*. 2022; 79:293-301.

6. Ansari Iari H, et al. In Vitro Comparison of the Effect of Three Types of Heat-Curing Acrylic Resins on the Amount of Formaldehyde and Monomer Release as well as Biocompatibility, *Advances in Materials Science and Engineering*. 2022; 8621666.
7. Azarpey A, et al., Bariatric surgery and secondary hyperparathyroidism; a mini-review, *Journal of Parathyroid Disease* 2023,11, e11238
8. Bauer P. R., et al. (2022). Plasma exchange in the intensive care unit: a narrative review. *Intensive Care Med*. 48, 1382–1396.
9. Beiranvandi F, et al., Investigation Of Medical Services In Patients With Diabetes And Cardio-Vascular Disease & High Blood Pressure In ICU With Radiological & Pathology Point: The Original Article, *Journal of Pharmaceutical Negative Results*, 2022; 4417-4425
10. Danesh H, et al., Pharmacological Evaluation of Covid 19 Vaccine in Acute and Chronic Inflammatory Neuropathies, *Journal of Medicinal and Chemical Sciences*, 2022; 5(4); 561-570
11. Effati, M., et al., Seven Year Prognosis and Surgical Techniques of Thoracic Esophageal Perforation Treatment, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, 2024, 3 (3), 672-681
12. Eghdam Zamiri, R., Mohammad Rahimi, M., Biomarkers Profile in Breast Carcinoma Presenting with Prostate Metastasis., *Eurasian Journal of Chemical, Medicinal and Petroleum Research*., 2024, 3 (2), 619-628
13. Fard MM, Fard AM, Investigation of drug release from a biodegradable biphasic polymer system, *Eurasian Journal of Science and Technology*, 2022; 2; 1-13
14. HSU, Chou-Yi, et al. Hydrazinosulfonylation of aryl electrophiles: a straightforward approach for the synthesis of aryl N-aminosulfonamides. *RSC advances*, 2023; 13.27: 18546-18560.
15. Imani R, Noroozi Seyed Hosseini H. Comment on "Factors affecting STEMI performance in six hospitals within one healthcare system". *Heart Lung*. 2023; 57:290. PMID: 35902305.
16. Jamali R, et al., *Journal of Pharmaceutical Negative Results*, 2022, 13(09)
17. Kareem HA, et al. Synthesis and Characterization of COFe₂O₄ Nanoparticles and Its Application in Removal of Reactive Violet 5 from Water. *J Nanostructures* 2022;12(3):521-528.
18. Keshmiri S, et al., *International Journal of Early Childhood Special Education*, 2022, 14 (1), 2960-2970
19. Kheradjoo, H., et al., Mesenchymal stem/stromal (MSCs)-derived exosome inhibits retinoblastoma Y-79 cell line proliferation and induces their apoptosis. *Nanomedicine Research Journal*, 2022; 7(3); 264-269.
20. Mahmoudi H, et al., *Nanomedicine Research Journal*, 2022, 7(3), 288-293,

21. Maroufi, P., Poulak, T., D-dimer Level Determination in the Diagnosis of Deep Vein Thrombosis Following Lower Extremity Fracture: a Cross-Sectional Study, *Eurasian Journal of Chemical, Medicinal and Petroleum Research.*, 2024, 3 (2), 651-658
22. Milanifard M, Weakness and Irritability, *GMJ Medicine*, 2021 5 (1), 391-395
23. Mirakhori F, et al., Diagnosis and Treatment Methods in Alzheimer's Patients Based on Modern Techniques: The Original Article, *Journal of Pharmaceutical Negative Results*, 2022, 13 (1), 1889-1907
24. Mirakhori F, et al., *Journal of Pharmaceutical Negative Results*, 2022, 1889-1907
25. Mohammadi G, I Seifi, SJ Barbin, E Zarei, R Tavakolimoghadam, *Tobacco Regulatory Science (TRS)*, 2022, 2064-2084
26. Motamedi T, et al., Investigating the Causes of Re-Laparotomy Surgery in the Field of Gastrointestinal Cancer in Patients Referred to Rasul Akram (PBUH) Educational and Therapeutic Complex During, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, 2023; 2 (1); 37-46
27. Nabiuni M, et al., Investigation of Types of Neuropathies in the Brain and Nerves, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, 2023; 2 (5); 1-15
28. Nabiuni M, et al., The Impact of Social Networks on Enhancing Safety and Efficacy Outcomes in Low-Dose Rituximab Treatment for Central Nervous System Demyelinating Diseases, *Interdisciplinary Journal of Virtual Learning in Medical Sciences*, 2023; 14 (3); 206-215
29. Noroozi Seyed Hosseini H, Imani R, Sumra B, et al. The effects of omega-3 and omega-6 fatty acids on the mental and physical health of children, adolescents, and adults. *Authorea*. 2023.
30. Otaghvar HA, et al., Investigating the Results of Amniocentesis in the Operating Room on Children's Acute Second Degree Burn Wounds in Patients Referred to Shahid Motahari Hospital, *Eurasian Journal of Chemical, Medicinal and Petroleum Research*, 2023; 2 (5); 32-44
31. Otaghvar HA, et al., Prevalence of Delirium and Its Related Factors in Burn Patients; a Systematic Review and Meta-Analysis, *Archives of Academic Emergency Medicine*, 2024; 12 (1); e7-e7
32. Parsa S, et al., BCL11B Is Involved in Stress-Induced Differentiation of Keratinocytes and Has A Potential Role in Psoriasis Pathogenesis, *Cell Journal (Yakhteh)*, 2023; 25 (5); 300
33. Rasouli F., Predicts factor in Polycystic Ovary Syndrome: An Evidence-based Study and Invitation from the Chief Editor., *Eurasian Journal of Chemical, Medicinal and Petroleum Research.*, 2024, 3 (2), 644-650
34. Rassam, M, Davoudi B., A Comprehensive Overview of Breast Cancer Surgery Strategies: Advances, Options, and Considerations., *Eurasian Journal of Chemical, Medicinal and Petroleum Research.*, 2024, 3 (2), 629-643

35. Rassam, M., Bahram, D., Relationship of CC49 with Outcomes after Colorectal Cancer Surgery., Eurasian Journal of Chemical, Medicinal and Petroleum Research., 2024, 2 (3), 610-618
36. Rustamzadeh A, et al., Horner syndrome: A new hypothesis for signaling pathway of enophthalmos sign, Current Journal of Neurology, 2023; 22 (3); 197
37. Seyedsalehi, AS., Rasouli, F., Do Adiponectin and Nesfatin Levels Predict Polycystic Ovary Syndrome (PCOS)? An Update Systematic Review, Eurasian Journal of Chemical, Medicinal and Petroleum Research., 2024, 3 (2), 659-671
38. Sharifi A, Dehghani A, Pain intensity after Esophagectomy in traumatic patients: Pioneering a New Era in Surgical Techniques, Eurasian Journal of Chemical, Medicinal and Petroleum Research, 2024; 3(1); 61-71
39. Sharifi A, Dehghani A, Short-term Outcomes of Gastrectomy for Patients with Gastric Cancer, Eurasian Journal of Chemical, Medicinal and Petroleum Research, 2024; 3 (1); 19-28
40. Sharifi A, Rousta F, Hypocalcemia: Why does this happen after thyroidectomy?, Eurasian Journal of Chemical, Medicinal and Petroleum Research, 2024; 3(1); 50-60
41. Sheikhy-Chaman M, Fazaeli AA, Darvishi A, Hadian M, Hamidi H. Social determinants of health with an emphasis on slum population. Social Determinants of Health. 2022;8(1):1-3.
42. Shiva Hoorzad AM, et al., Neuro Quantology, 2023, 21 (5), 316-324
43. Sidik MN, et al., In silico xanthine oxidase inhibitory activities of alkaloids isolated from Alphonsea sp. S Afr J Bot 2022;147; 820-825.
44. Taban M, et al., Maxillofacial Abnormalities and Surgical Stability After Changing the Angle of the Proximal Segment in Patients with Facial Asymmetry and Periodontal Problems, Seybold Report Journal (TSRJ), 2023; 18 (10); 1831-1851
45. Taban M, et al., Risk factors associated with implant sites prepared by orthodontic treatment: a systematic review, European Journal of Translational Myology, 2023; 33 (4)
46. Tomas, K. The physiology of wound healing. Annuals of Emergency Medicine, 2019;17(5). 1265-1288.
47. Yasrebinia S, Rezaei M, Artificial Intelligence for Predicting Neonatal Mortality in Post-Pregnancy: A Systematic Review, Eurasian Journal of Chemical, Medicinal and Petroleum Research, 2024; 3(1); 81-87
48. Zhang L, et al., Oncolytic viruses improve cancer immunotherapy by reprogramming solid tumor microenvironment, Medical Oncology, 2023; 41 (1); 8